


In the Claims:

The current status of all claims is listed below and supersedes all previous lists of claims.

Please amend claims 2, 4, 9, and 10 as follows:

1. (canceled).
2. (currently amended) A compound of formula (12):



wherein:

R^1 is hydrogen, C_{1-6} alkyl, or C_{3-6} cycloalkyl or a 4-10 membered aromatic mono or bicyclic ring containing 0 to 5 heteroatoms independently selected from nitrogen, oxygen or sulphur; wherein said C_{1-6} alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, carbamoyl, carboxy, C_{1-6} alkoxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl)₂amino, C_1 - C_6 alkylcarbonylamino, C_{1-6} alkylS(O)_a wherein a is 0-2, or C_{3-6} cycloalkyl or a 4-10 membered aromatic mono or bicyclic ring containing 0 to 5 heteroatoms independently selected from nitrogen, oxygen or sulphur; and wherein any said mono or bicyclic ring may be optionally substituted by one or two substituents selected from halo, hydroxy, C_{1-6} alkyl or C_{1-6} alkoxy;

R^2 and R^5 are independently hydrogen, a branched or unbranched C_{1-6} alkyl, or C_{3-6} cycloalkyl or a 4-10 membered aromatic mono or bicyclic ring containing 0 to 5 heteroatoms independently selected from nitrogen, oxygen or sulphur; wherein said C_{1-6} alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, cyano, carbamoyl, carboxy, C_{1-6} alkoxy, (a 4-10 membered aromatic mono or bicyclic ring containing 0 to 5 heteroatoms

independently selected from nitrogen, oxygen or sulphur) C₁₋₆alkoxy, (C₁₋₄alkyl)₃Si, N-(C₁₋₆alkyl)amino, N,N-(C₁₋₆alkyl)₂amino, C₁₋₆alkylS(O)_a, or C₃₋₆cycloalkyl, a 4-10 membered aromatic mono or bicyclic ring containing 0 to 5 heteroatoms independently selected from nitrogen, oxygen or sulphur or (a 4-10 membered aromatic mono or bicyclic ring containing 0 to 5 heteroatoms independently selected from nitrogen, oxygen or sulphur) C₁₋₆alkylS(O)_a, wherein a is 0-2; and wherein any said mono or bicyclic ring may be optionally substituted by one or two substituents selected from halo, hydroxy, C₁₋₆alkyl or C₁₋₆alkoxy;

R³ is hydrogen, C₁₋₆alkyl, halo, C₁₋₆alkoxy or C₁₋₆alkylS-

R⁴ is chlorine or fluorine;

R⁶ is ~~hydrogen~~, hydrogen or C₁₋₆alkyl, or (a 4-10 membered aromatic mono or bicyclic ring containing 0 to 5 heteroatoms independently selected from nitrogen, oxygen or sulphur) C₁₋₆alkyl;

wherein R⁵ and R² may form a ring with 2-7 carbon atoms and wherein R⁶ and R² may form a ring with 3-6 carbon atoms;

or a pharmaceutically acceptable salt, solvate, or a solvate of such a salt thereof.

3. (previously presented) A compound according to claim 2, wherein:

R¹ is hydrogen or phenyl.

4. (currently amended) A compound according to claim 2, wherein:

R² is hydrogen, a branched or unbranched C₁₋₆alkyl, or C₃₋₆cycloalkyl or a 4-10 membered aromatic mono or bicyclic ring containing 0 to 5 heteroatoms independently selected from nitrogen, oxygen or sulphur, wherein said C₁₋₆alkyl may be optionally substituted by one or more hydroxy, amino, acylamino, C₁₋₆alkylS(O)_a wherein a is 0-2, or C₃₋₆cycloalkyl or a 4-10 membered aromatic mono or bicyclic ring containing 0 to 5 heteroatoms independently selected from nitrogen, oxygen or sulphur; and wherein any said aromatic mono or bicyclic ring may be optionally substituted by hydroxy, C₁₋₆alkyl, alkoxy or cyano.

5. (previously presented) A compound according to claim 2, wherein:

R³ is hydrogen, C₁₋₂alkyl, halo or methoxy.

6. (previously presented) A compound according to claim 2, wherein:

R^3 is hydrogen, methyl, chlorine, fluorine, C_{1-6} alkyl-, or methoxy.

- 7-8. (canceled).

9. (currently amended) A compound according to claim 2, wherein:

R^6 is hydrogen, or C_{1-6} alkyl, ~~(a 4-10 membered aromatic mono or bicyclic ring containing 0 to 5 heteroatoms independently selected from nitrogen, oxygen or sulphur)-~~
 C_{1-6} alkyl or R^6 and R^2 form a ring with 3-6 carbon atoms.

10. (currently amended) A compound according to claim 2, wherein:

R^1 is hydrogen;

R^2 is a branched or unbranched C_{1-4} alkyl, optionally substituted by a C_{3-6} cycloalkyl, or
 C_{1-6} alkyl-S-, ~~a 4-10 membered aromatic mono or bicyclic ring containing 0 to 5 heteroatoms independently selected from nitrogen, oxygen or sulphur optionally substituted by hydroxy or~~
~~eyano~~, amino, N -(C_{1-6} alkyl)amino, or N,N -(C_{1-6} alkyl)₂amino ~~or (a 4-10 membered aromatic mono or bicyclic ring containing 0 to 5 heteroatoms independently selected from nitrogen,~~
~~oxygen or sulphur)-~~ C_{1-6} alkyl(S(O)_n wherein a is 0-2;

R^3 is halo;

R^5 is hydrogen or C_{1-6} alkyl; and

R^6 is hydrogen.

11. (previously presented) One or more compounds chosen from:

N -{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl- N^6 -acetyl-D-lysine;

1-(4-Fluorophenyl)-3-(R)-[2-(4-fluorophenyl)-2-hydroxyethylthio]-4-(R)-{4-[N -(N -[2-(phenyl)-1-(R)-(carboxy)ethyl]carbamoylmethyl)carbamoylmethoxy]phenyl]azetidin-2-one;

N -{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-valine;

N -{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-

oxoazetidin-2-yl)phenoxy]acetyl} glycyL-D-tyrosine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl} glycyL-D-proline;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl} glycyL-D-lysine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-hydroxy-2-(4-methoxyphenyl)ethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl} glycyL-D-valine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl} glycyL-2-butylnorleucine;

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl} glycyL-*S*-methyl-L-cysteine;

N-{[4-((2*R*,3*R*)-1-(4-chlorophenyl)-3-{[2-(4-chlorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl} glycyL-3-cyclohexyl-D-alanine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl} glycyL-3-cyclohexyl-D-alanine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl} glycyL-4-methylleucine;

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-L-alanyl-D-valine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-hydroxy-2-(4-methylphenyl)ethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl} glycyL-D-valine;

N-{[4-((2*R*,3*R*)-1-(4-chlorophenyl)-3-{[2-(4-chlorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl} glycyL-D-valine;

N-{[4-((2*R*,3*R*)-1-(4-chlorophenyl)-3-{[2-(4-chlorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl} glycyL-3-methyl-D-valine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl} glycyL-3-(2-naphthyl)-D-alanine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl} glycyL-3-methyl-D-valine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-

oxoazetidin-2-yl)phenoxy]acetyl}glycyl-(3*R*,4*S*,5*R*)-3,4,5,6-tetrahydroxy-D-norleucine;

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-*N*,2-dimethylalanine;

N-{[4-[(2*R*,3*R*)-1-(4-Fluorophenyl)-3-{2-hydroxy-2-[4-(methylthio)phenyl]ethyl}thio]-4-oxoazetidin-2-yl]phenoxy]acetyl}glycyl-3-methyl-D-valine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-*S*-(4-methylbenzyl)-D-cysteine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-*S*-(*tert*-butyl)-D-cysteine; and

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-*b*,*b*-dimethyl-D-phenylalanine.

12. (canceled).


13. (previously presented) A method of treating a hyperlipidemic condition comprising the administration of an effective amount of a compound according to claim 2 to a mammal in need thereof.

14. (previously presented) A method of treating atherosclerosis comprising the administration of an effective amount of a compound according to claim 2 to a mammal in need thereof.

15-16. (canceled).

17. (previously presented) A pharmaceutical formulation comprising a compound according to claim 2 in admixture with a pharmaceutically acceptable adjuvant, diluent and/or carrier.

18. (previously presented) A combination of a compound according to formula (I)



wherein:

R^1 is hydrogen, C_{1-6} alkyl, C_{3-6} cycloalkyl or aryl; wherein said C_{1-6} alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, carbamoyl, carboxy, C_{1-6} alkoxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino, C_1 - C_6 alkylcarbonylamino, C_{1-6} alkylS(O) $_a$ wherein a is 0-2, C_{3-6} cycloalkyl or aryl; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C_{1-6} alkyl or C_{1-6} alkoxy;

R^2 and R^5 are independently hydrogen, a branched or unbranched C_{1-6} alkyl, C_{3-6} cycloalkyl or aryl; wherein said C_{1-6} alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, cyano, carbamoyl, carboxy, C_{1-6} alkoxy, aryl C_{1-6} alkoxy, $(C_1-C_4alkyl)_3Si$, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkylS(O) $_a$, C_{3-6} cycloalkyl, aryl or aryl C_{1-6} alkylS(O) $_a$, wherein a is 0-2; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C_{1-6} alkyl or C_{1-6} alkoxy;


R^3 is hydrogen, alkyl, halo, C_{1-6} alkoxy or C_{1-6} alkylS-;

R^4 is hydrogen, C_{1-6} alkyl, halo or C_{1-6} alkoxy;

R^6 is hydrogen, C_{1-6} alkyl, or aryl C_{1-6} alkyl;

wherein R^5 and R^2 may form a ring with 2-7 carbon atoms and wherein R^6 and R^2 may form a ring with 3-6 carbon atoms;

or according to formula (I2)



wherein:

R¹ is hydrogen, C₁₋₆alkyl, C₃₋₆cycloalkyl or aryl; wherein said C₁₋₆alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, carbamoyl, carboxy, C₁₋₆alkoxy, *N*-(C₁₋₆alkyl)amino, *N,N*-(C₁₋₆alkyl)₂amino, C₁-C₆alkylcarbonylamino, C₁₋₆alkylS(O)_a wherein a is 0-2, C₃₋₆cycloalkyl or aryl; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C₁₋₆alkyl or C₁₋₆alkoxy;

R² and R⁵ are independently hydrogen, a branched or unbranched C₁₋₆alkyl, C₃₋₆cycloalkyl or aryl; wherein said C₁₋₆alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, cyano, carbamoyl, carboxy, C₁₋₆alkoxy, aryl C₁₋₆alkoxy, (C₁-C₄alkyl)₃Si, *N*-(C₁₋₆alkyl)amino, *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkylS(O)_a, C₃₋₆cycloalkyl, aryl or aryl C₁₋₆alkylS(O)_a, wherein a is 0-2; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C₁₋₆alkyl or C₁₋₆alkoxy;

R³ is hydrogen, alkyl, halo, C₁₋₆alkoxy or C₁₋₆alkylS-;


R⁴ is hydrogen, C₁₋₆alkyl, halo or C₁₋₆alkoxy;

R⁶ is hydrogen, C₁₋₆alkyl, or arylC₁₋₆alkyl;

wherein R⁵ and R² may form a ring with 2-7 carbon atoms and wherein R⁶ and R² may form a ring with 3-6 carbon atoms;

with a PPAR alpha and/or gamma agonist.

19. (previously presented) A combination of a compound according to formula (I)



wherein:

R^1 is hydrogen, C_{1-6} alkyl, C_{3-6} cycloalkyl or aryl; wherein said C_{1-6} alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, carbamoyl, carboxy, C_{1-6} alkoxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino, C_1 - C_6 alkylcarbonylamino, C_{1-6} alkylS(O) $_a$ wherein a is 0-2, C_{3-6} cycloalkyl or aryl; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C_{1-6} alkyl or C_{1-6} alkoxy;

R^2 and R^5 are independently hydrogen, a branched or unbranched C_{1-6} alkyl, C_{3-6} cycloalkyl or aryl; wherein said C_{1-6} alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, cyano, carbamoyl, carboxy, C_{1-6} alkoxy, aryl C_{1-6} alkoxy, $(C_1-C_4alkyl)_3Si$, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkylS(O) $_a$, C_{3-6} cycloalkyl, aryl or aryl C_{1-6} alkylS(O) $_a$, wherein a is 0-2; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C_{1-6} alkyl or C_{1-6} alkoxy;


R^3 is hydrogen, alkyl, halo, C_{1-6} alkoxy or C_{1-6} alkylS-;

R^4 is hydrogen, C_{1-6} alkyl, halo or C_{1-6} alkoxy;

R^6 is hydrogen, C_{1-6} alkyl, or aryl C_{1-6} alkyl;

wherein R^5 and R^2 may form a ring with 2-7 carbon atoms and wherein R^6 and R^2 may form a ring with 3-6 carbon atoms;

or according to formula (I2)



wherein:

R^1 is hydrogen, C_{1-6} alkyl, C_{3-6} cycloalkyl or aryl; wherein said C_{1-6} alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, carbamoyl, carboxy, C_{1-6} alkoxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino, C_1 - C_6 alkylcarbonylamino, C_{1-6} alkylS(O) $_a$ wherein a is 0-2, C_{3-6} cycloalkyl or aryl; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C_{1-6} alkyl or C_{1-6} alkoxy;

R^2 and R^5 are independently hydrogen, a branched or unbranched C_{1-6} alkyl, C_{3-6} cycloalkyl or aryl; wherein said C_{1-6} alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, cyano, carbamoyl, carboxy, C_{1-6} alkoxy, aryl C_{1-6} alkoxy, $(C_1-C_4alkyl)_3Si$, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkylS(O) $_a$, C_{3-6} cycloalkyl, aryl or aryl C_{1-6} alkylS(O) $_a$, wherein a is 0-2; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C_{1-6} alkyl or C_{1-6} alkoxy;

R^3 is hydrogen, alkyl, halo, C_{1-6} alkoxy or C_{1-6} alkylS-;

R^4 is hydrogen, C_{1-6} alkyl, halo or C_{1-6} alkoxy;

R^6 is hydrogen, C_{1-6} alkyl, or aryl C_{1-6} alkyl;

wherein R^5 and R^2 may form a ring with 2-7 carbon atoms and wherein R^6 and R^2 may form a ring with 3-6 carbon atoms;

with an HMG Co-A reductase inhibitor.

20-28. (canceled).

29. (previously presented) A combination of a compound according to claim 2 with a PPAR alpha and/or gamma agonist.

30. (previously presented) A combination of a compound according to claim 2 with an HMG Co-A reductase inhibitor.